

SELECTED PUBLICATIONS

GERHARD SATTLER, MD



- Sattler G, Eichner S. Complications of liposuction. *Hautarzt*. 2013;64(3):171-9.
- Krueger N, Luebberding S, Sattler G, William Hanke C W, Alexiades-Armenakas M, Sadick N. The history of aesthetic medicine and surgery. *J Drugs Dermatol*. 2013; 12(7):737-42.
- Sattler G et al. Recommendations for the best possible use of botulinum neurotoxin type a (Speywood units) for aesthetic applications. *J Drugs Dermatol*. 2012;12(1):80-4.
- Sattler G. The tower technique and vertical supraperiosteal depot technique: novel vertical injection techniques for volume-efficient subcutaneous tissue support and volumetric augmentation. *J Drugs Dermatol*. 2012;11(8):s45-7.
- Sattler G et al. Anatomic guidelines for augmentation of the cheek and infraorbital hollow. *Dermatol Surg*. 2012 Jul;38(7 Pt 2):1223-33.
- Sattler G et al. Validated assessment scale for neck volume. *Dermatol Surg*. 2012;38(2 Spec No.):343-50.
- Sattler G. What does the future hold for botulinum neuromodulators in facial aesthetics? *J Eur Acad Dermatol Venereol*. 2012;26(3):394-5.
- Rzany B, Cartier H, Kestermont P, Trevidic P, Sattler G, Kerrouche N, Dhuin JC. Correction of tear troughs and periorbital lines with a range of customized hyaluronic acid fillers. *J Drugs Dermatol*. 2012; 11(S1):27-34.
- Rzany B, Cartier H, Kestemont, P, Trevidic P, Sattler G, Kerrouche N, Dhuin, J-C, May Ma Y. Full-face rejuvenation using a range of hyaluronic acid fillers. Efficacy, safety, and patient satisfaction over 6 months. *Dermatol Surg*. 2012; 38(7 Pt 2):1153-61.
- Cartier H, Trevidic P, Rzany B, Sattler G, Kestemont P, Kerrouche N, Dhuin JC. Perioral rejuvenation with a range of customized hyaluronic acid fillers. Efficacy and safety over six months with a specific focus on the Lips. *J Drugs Dermatol*. 2012; 11(S1):15-26.
- Bartus C, Sattler G, Hanke C. The Tower Technique. A Novel Technique for the Injection of Hyaluronic Acid Fillers. *J Drugs Dermatol*. 2011; 10(11):1277-80.
- Sattler G et al. Noninferiority of incobotulinumtoxinA, free from complexing proteins, compared with another botulinum toxin type A in the treatment of glabellar frown lines. *Dermatol Surg*. 2010;36 Suppl 4:2146-54.
- Sattler G et al. Double-blind, randomized, intraindividual comparison study of the efficacy of prilocaine and lidocaine in tumescent local anesthesia. *Dermatology*. 2010;221(3):248-52.
- Sattler G. Current and future botulinum neurotoxin type A preparations in aesthetics: a literature review. *J Drugs Dermatol*. 2010;9(9):1065-71.
- Sattler G et al. Improvement in skin elasticity in the treatment of cellulite and connective tissue weakness by means of extracorporeal pulse activation therapy. *Aesthet Surg J*. 2008;28(5):538-44.
- Carruthers A, Carruthers J, Hardas B, Kaur M, Görtelmeyer R, Jones D, Rzany B, Cohen J, Kersch M, Flynn TC, Maas C, Sattler G, Gebauer A, Pooth R, McClure K, Simone-Korbel U, Buchner L. A validated hand grading scale. *Derm Surg*. 2008; 11(S2): 179-83.

Christ C, Brenke R, Sattler G, Siems W, Novak P, Daser A. Improvement in skin elasticity in the treatment of cellulite and connective tissue weakness by means of extracorporeal pulse activation therapy: EPAT®. *Aesthet Surg J*. 2008; 28(59): 538-44.

Sattler et al. Consensus recommendations on the use of botulinum toxin type A in aesthetic medicine. *J Dtsch Dermatol Ges*. 2007 Feb;5 Suppl 1: S1-S29.

Sattler G. Advances in liposuction and fat transfer. *Dermatol Nurs*. 2005; 17(2): 133-9.

Grossmann M, Sattler G, Pistner H, Oertel R, Richter K, Schinzel S, Jacobs LD. Pharmacokinetics of articaine hydrochloride in tumescent local anesthesia for liposuction. *J Clin Pharmacol*. 2004;44(11):1282-9.

Sattler G, Bergfeld D, Sommer B. Liposuktion. *Hautarzt*. 2004; 55(7): 599-604.

Sommer B, Bergfeld D, Sattler G. LipoRepair. Behandlungskonzept bei Fettgewebsdeformitäten *Hautarzt*. 2004; 55(7): 605-10.

Wörle B, Hanke CW, Sattler G. Poly-L-Lactic Acid. A temporary filler for soft tissue augmentation. *J Drugs Dermatol*. 2004; 3(4): 385-389.

Sommer B, Zschocke I, Bergfeld D, Sattler G, Augustin M. Satisfaction of patients after treatment with botulinum toxin for dynamic facial lines. *Dermatol Surg*. 2003;29(5):456-60.

Sattler G. Patient management and treatment technique. *Clinics Dermatol*. 2003; 21(6): 485-7.

Pröbstle TM, Bethge S, Barnstedt S, Kargl A, Knop J, Sattler. Subfascial endoscopic perforator surgery with tumescent local anesthesia. *Dermatol Surg*. 2002;28(8):689-93.

Sommer B, Sattler G. Botulinum toxin in aesthetic medicine. *J Cosmet Dermatol*. 2002; 1(2): 103-4.

Augustin M, Zschocke I, Godau N, Buske-Kirschbaum A, Peschen M, Sommer B, Sattler G. Skin surgery under local anesthesia leads to stress-induced alterations of psychological, physical, and immune functions. *Derm Surg*. 2001; 25(11): 868-71.

Sommer B, Sattler G. Current concepts of fat graft survival. Histology of aspirated adipose tissue and review of the literature. *Dermatol Surg*. 2000. 26(12): 1159-66.

Sattler G. Subfascial endoscopic perforator surgery. *Curr Probl Dermatol*. 1999;27:190-4.